DR EYAS MEDANAT

Dr. Eyas Medanat
Unit 303, 3rd Floor, Block B, Ibn Sina Building 27, Dubai Healthcare City, Dubai, UAE
+971 50 108 8378 · info@eyas.doctor

GUIDES · BREAST AUGMENTATION

Breast Augmentation (Implants)

Breast augmentation places an implant behind the breast tissue or chest muscle to add volume and shape. Implants take around 3 months to settle into their final position — early on they sit higher and feel firm, which is expected.



1 At a glance

PROCEDURE TIME	ANAESTHESIA	GOING HOME	GARMENT / DRESSING
About 1–1.5 hours	General anaesthesia	Home the same day or next morning	Surgical support bra 24/7 for 6 weeks (no underwire)

STITCHES	BACK TO WORK	EXERCISE
Usually dissolvable, under skin tape	1–2 weeks (desk work)	Walking from day 1; lower-body gym at 4 weeks; chest and arms at 6–8 weeks

2 Before your surgery

Specific to this procedure

- Buy two surgical support bras in the size we advise (front-fastening is easiest).
- If you are over 40 or have a family history of breast cancer, a recent breast scan may be needed first.
- Prepare things at waist-to-shoulder height at home so you are not reaching up for the first weeks.

Preparing for surgery

2–4 WEEKS BEFORE

- Stop smoking and all nicotine — cigarettes, vapes, shisha, patches and gum — at least 4 weeks before and 4 weeks after surgery. Nicotine narrows the small blood vessels that heal your wounds and is the commonest cause of wound breakdown.
- Complete the blood tests, ECG or other checks we request, and bring any recent results you already have.
- Tell us about every medicine, supplement and herbal product you take, any allergies, and any previous anaesthetic problems.
- If you take the contraceptive pill, hormone therapy, or weight-loss injections (such as Ozempic, Wegovy or Mounjaro), tell us early — some need to be paused before anaesthesia, and we will tell you exactly when.
- Arrange time off work and help at home for the first days, especially if you look after young children.

10 DAYS BEFORE

- Stop aspirin, ibuprofen and other anti-inflammatory painkillers, and supplements that thin the blood: fish oil / omega-3, vitamin E, ginkgo, garlic, ginseng and turmeric. Paracetamol is fine.
- Do not stop any prescribed blood thinner (warfarin, apixaban, clopidogrel and similar) on your own — we will plan this with you and your doctor.
- Avoid alcohol from one week before surgery.
- Let us know straight away if you develop a cold, fever, cough, cold sore or any skin infection — surgery may need to move to keep you safe.

THE DAY BEFORE

- Nothing to eat for 6 hours before your admission time. You may drink plain water until 2 hours before, unless we give you different times.
- Take your usual morning medicines with a sip of water only if we have told you to.
- Prepare your recovery space: extra pillows, loose front-opening clothes, water bottle, phone charger within reach.
- Arrange an adult to drive you home and stay with you for the first 24 hours. You must not drive or take a taxi alone after anaesthesia or sedation.

ON THE DAY

- Shower and wash your hair. Do not apply body lotion, make-up, false lashes, nail polish, fake tan, perfume or deodorant.
- Remove jewellery and piercings; leave valuables at home. Wear glasses rather than contact lenses.
- Wear loose, comfortable clothes that open at the front and flat shoes.
- Bring your ID, your list of medicines, and your compression garment if we asked you to buy one.

3 After your surgery

Your medicines

- Augmentin (antibiotic): 1 tablet twice a day, with food, for 7 days. Finish the full course even if you feel well. Tell us before surgery if you are allergic to penicillin.
- Reparil (for swelling and bruising): 2 tablets three times a day for 7–14 days, as we tell you.
- Brufen (painkiller): take it only when you need it, with food. If you cannot take anti-inflammatory painkillers — for example because of stomach ulcers, kidney problems or asthma — tell us and we will give you an alternative.
- Fusidic acid cream (antibiotic cream): apply a thin layer on the wounds twice a day, after showering, until they have healed.

First week

- Rest with your back raised at 30–45 degrees; sleep on your back for 4–6 weeks.
- Walk around the house every couple of hours to protect your circulation.
- Keep your elbows close to your body — no reaching above your shoulders, pushing up from the bed with your arms, or lifting more than 2–3 kg.
- Tightness across the chest and a burning feeling in the nipples are common as the tissues stretch.

Wounds and the bra

- Wear the support bra (and band, if we gave you one) day and night for 6 weeks, taking it off only to shower.
- Shower from 48 hours; leave the skin tapes on, pat them dry. We will change or remove them at your check.
- No baths, swimming pools, sea or saunas for 4 weeks.
- From 3 weeks, once the wounds have healed, start silicone scar gel or tape.

Longer term

- No underwire bras for 3 months.

- Keep your implant card safe — it has the make, size and serial number.
- Tell anyone doing a mammogram or ultrasound in future that you have implants.
- Breast implants are not lifetime devices; we recommend a check with us every year or if you notice any change.

4 Your recovery, week by week

Day 1–3
Most tightness and discomfort. Up and walking.

Week 1
First check. Back to desk work, driving once you can brake hard and turn without pain.

Weeks 4–6
Lower-body exercise; bra continues to 6 weeks.

Weeks 6–8
Upper-body exercise and normal sports bras.

Month 3
Implants have dropped and softened; final shape.

5 Follow-up visits

- 4 days after your procedure
- 2 weeks after your procedure
- 4 weeks after your procedure
- 6 weeks after your procedure

Call us straight away if you notice

⚠ One breast becoming much larger, harder or more painful than the other.

⚠ Swelling, redness or fluid appearing weeks or months after surgery.

⚠ A temperature of 38 °C or higher, shivering, or feeling generally unwell.

⚠ Bleeding that soaks through the dressing, or does not stop after 10 minutes of firm pressure.

⚠ Swelling that is sudden, tense and painful, or clearly larger on one side than the other.

⚠ Redness that is spreading, increasing heat, pus, or a bad smell from a wound.

⚠ Pain that is getting worse rather than better, or not controlled by the medicine we gave you.

⚠ Pain, swelling or redness in a calf, or new breathlessness or chest pain — call 998 first.

⊕ **Chest pain, difficulty breathing, or heavy bleeding that will not stop: call 998 (ambulance) or go to the nearest emergency department, then let us know.**

Questions while you recover — Message or call the clinic any time — no question is too small, and it is always better to ask early. **+971 50 108 8378** (WhatsApp) · info@eyas.doctor

---

This guide gives general advice. Dr. Eyas may give you instructions made just for you — at your consultation or when you go home after surgery. If his instructions are different from this guide, follow his instructions. If you are not sure, message us. Last reviewed: November 2026.